

(9)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

**0 315 349
A1**

(12)

EUROPEAN PATENT APPLICATION

(2) Application number: **88309996.2**

(51) Int. Cl. 4: **A61K 47/00**

(22) Date of filing: **24.10.88**

Claims for the following Contracting States: GR
+ ES.

(30) Priority: **23.10.87 IL 84252**

(43) Date of publication of application:
10.05.89 Bulletin 89/19

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **YISSUM RESEARCH DEVELOPMENT
COMPANY OF THE HEBREW UNIVERSITY OF
JERUSALEM**
46 Jabotinsky Street
Jerusalem, 92 182(IL)

(72) Inventor: **Yedgar, Saul**
1 Teib Street
Jerusalem(IL)
Inventor: **Dagan, Arie**
13 harlap Street
Jerusalem(IL)

(74) Representative: **Sheard, Andrew Gregory et al**
Kilburn & Strode 30, John Street
London WC1N 2DD(GB)

(54) **Phospholipase A₂ inhibiting compositions and their use.**

(57) Inhibitors of phospholipase A₂ activity at the cell-surface membrane whose molecular structure comprises a cell-permeable PLA₂-inhibitor moiety covalently bonded directly or indirectly to a physiologically acceptable carrier moiety which is effective to inhibit cell internalisation of the cell-permeable PLA₂-inhibitor moiety, with the proviso that phosphatidylserine is not bonded indirectly via divalent dodecanedioyl to dextrane hydrazide.

EP 0 315 349 A1

PHOSPHOLIPASE A₂ INHIBITING COMPOSITIONS AND THEIR USE

This invention relates to novel compositions of matter which are inhibitors of the enzyme phospholipase A₂ (PLA₂) at the cell membrane, to methods for their production and to methods for their use to treat PLA₂-related abnormal conditions.

Phospholipase A₂ (PLA₂, EC 3.1.1.4), which hydrolyzes phospholipids at the $\rho\eta$ -2 position to produce a fatty acid and a lysophospholipid, is present in almost all cell types and subcellular fractions. See Van den Bosch, H. (1982) "Phospholipids" (Hawthorne, N.J. & Ansell, G.D., Editors), pp. 313-357 (Elsevier Pub., Amsterdam, N.Y.). The activity of this enzyme has been correlated with various cell functions, particularly with secretory processes such as exocytosis and eicosanoid production (prostaglandins, thromboxanes and leukotrienes). See Waite M. (1987). "Phospholipases", Plenum Press, N.Y. Accordingly, PLA₂-inhibitors were proposed for treatment of diseases which are associated with enhanced cellular secretion, such as allergy (secretion of histamine) inflammation (secretion of lysosomal enzymes and peroxides) or thrombosis (secretion of thromboxane).

It is accepted that the enzyme involved in these processes is located in the cell-surface membrane (the plasma membrane), and regulation of the membrane-enzyme's activity is required for regulation of cellular secretion. See Blackwell, G.J. and Flower, R.J. (1983). Br. Med. Bull. 39, 260-264. A number of PLA₂ inhibitors have been considered for treatment of oversecretion-related pathological states. Among them are the lipocortin-like proteins which appear in mammalian tissues and assumed to be induced by glucocorticoids (although this has not been unequivocally proven). However, the prolonged administration of steroids has many side effects and, in general, is undesirable. The lipocortin-like proteins can be provided exogenously and might affect the cell membrane PLA₂ activity and cellular secretion. However, since these substances are calcium-binding proteins they undesirably interfere with many other cellular functions. See: Crompton et al. (1988) Cell, 55, 1-3.

Other inhibitors, including N-derivatives of phosphatidylserine (see Martina Lagunoff (1982) Biochemistry 21, 1254-1260) are synthetic or natural products of smaller molecular weight which can be administered exogeneously. However, they are also internalized by the cell, interfere with the vital lipid metabolism and are therefore cytotoxic. An extra-cellular inhibitor of PLA₂ which affects the enzyme at the cell membrane surface but does not penetrate into the cell would, therefore, be very desirable. The PLA₂ inhibitors of this invention fulfill these requirements.

In a paper published in FEBS Lett. (1986) 200 (1), pp. 165-8, which we coauthored with Nurit Reisfeld, we report the synthesis of a cell-impermeable inhibitor of phospholipase A₂, by acylating the amino group of phosphatidylserine (PS) with dodecanedioic acid and linking the resulting free carboxyl group to a high molecular weight (70,000) dextran-hydrazide. This inhibitor incorporates into lipid membranes and is capable of blocking the hydrolysis of membrane phospholipids by snake venom as well as by cell membrane PLA₂.

This specific PLA₂ inhibitor, although cell-impermeable, has been abandoned since dextran-hydrazide is an undesirable moiety from a physiological point of view, and we have since found that linkage of PLA₂ inhibiting moieties to other carrier (polymer) moieties produces cell-impermeable PLA₂-inhibitors with markedly better properties. These preparations are capable of inhibiting cellular secretion and eicosanoid production, as well as membrane-PLA₂ activity in intact cells without impairing the cell viability.

In a first aspect, this invention relates to chemical compounds which are inhibitors of the enzyme phospholipase A₂ (PLA₂) and whose molecular structures comprise a cell-permeable PLA₂-inhibitor moiety which interacts with the enzyme at the cell plasma membrane, covalently bonded preferably directly or indirectly to a physiologically acceptable carrier moiety which is effective to inhibit the cellular internalisation of the cell permeable PLA₂-inhibiting moiety, with the proviso that phosphatidylserine is not bonded indirectly via divalent dodecanedioyl to dextran-hydrazide. Preferably the carrier moiety is not dextran-hydrazide.

In a second aspect, this invention relates to a method of ameliorating the adverse symptoms exhibited by a living being suffering from a PLA₂-related pathological condition, which comprises administering thereto an amount of a compound of this invention effective to ameliorate those symptoms, or a PLA₂-inhibiting amount per unit dosage of a compound of the first aspect.

In a third aspect, the invention relates to processes for the conversion of a cell-permeable PLA₂ inhibitor into impermeable derivatives which are inhibitors of the cell-membrane PLA₂ activity.

A fourth aspect relates to a pharmaceutical composition comprising a compound capable of inhibiting the enzyme phospholipase A₂ (PLA₂) comprising a cell-permeable PLA₂-inhibitor moiety covalently bonded

directly or indirectly to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, with the proviso that phosphatidylserine is not bonded indirectly via a divalent didodecanedioyl group to dextran-hydrazide and a pharmaceutically acceptable carrier.

A fifth aspect extends to the use of a compound capable of inhibiting the enzyme phospholipase A₂ - (PLA₂) comprising a cell-permeable PLA₂-inhibitor moiety covalently bonded directly or indirectly to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, in the preparation of a PLA₂ inhibiting agent.

A sixth aspect of the invention encompasses a process for the preparation of a compound capable of inhibiting the enzyme phospholipase A₂ (PLA₂), the process comprising covalently bonding directly or indirectly a cell-permeable PLA₂-inhibitor moiety to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, with the proviso that phosphatidylserine is not bonded indirectly via divalent didodecanedioyl to dextran-hydrazide. Preferred features for one aspect are for another aspect *mutatis mutandis*.

The PLA₂ inhibitors of this invention comprise in their molecular structure cell-permeable PLA₂ inhibitors, many of which are known in the prior art. The term "moiety" means a chemical entity otherwise corresponding to a chemical compound, which has a valence satisfied by a covalent bond. Typically, the PLA₂-inhibitor moiety will be identical in structure to a chemical compound having PLA₂-inhibiting activity, except at the point at which that moiety is covalently bonded to the carrier portion of the molecule of the PLA₂ inhibitor of this invention, e.g., it differs therefrom by having a valence which replaces a hydrogen atom.

PLA₂ inhibitors can be classified into the following types:


I. Phosphatidylethanolamine (PE) and its analogues, such as distearoyl-PE (which gave the best results). Natural PEs from various sources, semisynthetic PEs, synthetic natural and artifactual (novel, unnatural) PEs and their isomers. Any of the above-mentioned compounds linked through the amino group of the ethanolamine by a covalent bond.

II. N-methyl-PE derivatives and their analogues, linked through the amino group of the N-methyl-PE by a covalent bond.

III. N,N-dimethyl-PE derivatives and their analogues linked through the amino group of the N,N-dimethyl-PE by a covalent bond.


IV. Phosphatidylserine (PS) and its analogues, such as Natural PSs from various sources, semisynthetic PSs, synthetic natural and artifactual PSs and their isomers. Any of the above-mentioned compounds linked through the PS amino group by a covalent bond.

V. Glycerol ether, amine, amide, thioether, ester and thioester derivatives of the general formula




wherein X is -O-, -S-,

-O- $\overset{\text{O}}{\parallel}\text{C}$ -, -O- $\overset{\text{S}}{\parallel}\text{C}$ -, -NH-, -NH- $\overset{\text{O}}{\parallel}\text{C}$ -; R₁ is alkyl, alkyl-COOH, alkylamine e.g. where alkyl means C₁₋₁₀ alkyl; R₂ is alkyl, alkyl-COOH, alkylamine e.g. where alkyl means C₁₋₁₅ alkyl; R₃ is alkyl, phosphate, phosphorylcholine, phosphorylserine, phosphorylglycerol, phosphorylinositol, etc., linked to a carrier via one (or more) of the R groups. Best results were obtained with the following compound:





The term "alkyl" in this specification may denote a C₁₋₁₄ alkyl group unless otherwise stated.

VI. Ethylene glycol derivatives of the general formula




linked to a carrier via one of the R groups, wherein R₁ is alkyl,

- $\overset{\text{O}}{\parallel}{\text{C}}$ -alkyl e.g. C₁₋₁₁ alkyl, phosphorylcholine, phosphorylserine, phosphorylethanolamine, phosphorylglycerol, phosphonylinositol, phosphate, etc.; R₂ is alkyl, alkyl-COOH, alkylamine; X is -O-, -S-,



Best results were obtained with the following:



VII. Aminopiperazine and its derivatives.

VIII. Manoalide and its derivatives, synthetic and natural, e.g., manoalogue.

IX. Arachidonic acid and its derivatives, natural and synthetic.

X. p-methoxyphenethylamine, its analogues and derivatives.

XI. Sphingosines, their analogues and derivatives.

XII. Phenacylbromides.

Examples of cell-penetrating PLA₂ inhibitors which can be employed as the PLA₂-inhibitor moiety of the compounds of this invention are aminophospholipids, (e.g., phosphatidylethanolamine, phosphatidylserine), mepacrine, local anesthetics, e.g., chlorpromazine, procaine, indomethacin, sulfated analogs of indomethacin, bromophenacyl bromide, p-methoxyphenethylamine, imipramine, propanolol, phenothiazines, quinacrine, dibucaine, tetracaine, lidocaine, 1-amino-4-octylpiperazine, 1,7-bis(p-aminophenoxy)heptane, tripelennamine, amantadine and phentermine, manoalide, manoalogue, sphingosine and derivatives of each of the above which possess PLA₂ inhibiting activity.

The primary role of the carrier moiety is to increase the size (molecular volume) of the PLA₂ inhibitor forming the PLA₂-inhibitor moiety of the composition of this invention sufficient to render the latter cell-impermeable. When the starting carrier molecule has a substituent which is or can be rendered reactive to

a substituent on the starting PLA₂-inhibitor compound, the carrier molecule can be linked directly to the PLA₂-inhibitor molecule. When it does not, a bifunctional linking starting material can be used to link the two molecules together indirectly.

Examples of carrier moieties which can be employed to eliminate the cell-penetrating ability of the PLA₂-inhibitor portion (moiety) of this invention are physiologically acceptable polymers, including water-dispersible or -soluble polymers of various molecular weights and diverse chemical types, mainly plasma expanders and food and drug additives, including "Hemacell" (degraded gelatin polypeptide crosslinked via urea bridges, produced by "Behring") "hydroxyethyl-starch" (HES) polyamino acids, hydrocarbon polymers (e.g., polyethylene), polystyrenes, polyesters, polyamides, polyethylene oxides (e.g., polyethyleneglycols), polyvinylpyrrolidones, polysaccharides, soluble cellulose derivatives (e.g., methylcellulose, carboxymethyl cellulose), alginates, assimilable gums (e.g., xanthan gum, stractan), peptides, injectable blood proteins (e.g., serum albumin), cyclodextrin, and derivatives thereof.

The carriers can have a wide range of molecular weight, e.g., above 50,000 (up to a few hundred thousands e.g. 300,000) when it is desirable to retain the DLA₂ inhibitor in the vascular system and below 50,000 when targeting to extravascular systems is desirable. The sole limitation on the molecular weight and the chemical structure of the carrier moiety is that it does not destroy the PLA₂-inhibiting activity of the PLA₂-inhibiting moiety and does not promote cellular uptake of the PLA₂ inhibitor.

Examples of suitable divalent groups forming the optional bridging group are straight- or branched-chain alkylene, e.g., of 2 or more, preferably 4 to 18 carbon atoms, -CO-alkylene-CO-, -NH-alkylene-NH-, -CO-alkylene-NH-, cycloalkylene, wherein alkylene in each instance, is straight or branched chain and contains 2 or more, preferably 2 to 18 carbon atoms in the chain, {O-ethylene}_X-O- and {O-CH(CH₃)-CH₂}_X-O- wherein X is an integer of 1 or more.

Illustrative of compounds of this invention are those in which a PLA₂-inhibitor moiety listed below is linked directly or indirectly through a bridging moiety listed below, to a carrier moiety listed below.

	<u>PLA₂-Inhibitor Moiety</u>	<u>Bridging Moiety</u>	<u>Carrier Moiety (MW)</u>
5	N-derivatized-phosphatidyl- -ethanolamines	aminoacids	"Hemacells"
	N-derivatized-phosphatidyl- -serine	diaminoalkyls	carboxymethyl- cellulose
10	Glycerol-ethers and thioethers	diacylcarboxylic acids	methylcelluloses
	Glycerol-amines and amides	dialcohols	alginates
15	Glycerol esters and thioesters	dimercaptans	hydroxyethyl starch
	Ethylene glycol derivatives	dialdehydes	polyethylenes
20	Retinoides	formylcarboxylic acids	dextrans
	Manoalide, mancaloge and their derivatives		albumins
25	Arachidonic acid		cyclodextrins
	Methoxyphenetylamines		polyamino acids
30	Sphingosines		
	Phenacylbromides		
35	Aminopiperazines		

In addition to the compounds of the Examples, further illustrative compounds of this invention are set forth in the table below.


CMC-PS; CMS-PE;
 Hemacell-PS; Hemacell-PE (Hemacell M.W. = 35,000);
 Alginic-PS; Alginic-PE;
 Manoalide-Hemacell;
 Manoalogue-dextran (MM 40,000);
 p-bromophenacylbromide-Hemacell;
 glycerolether-Hemacell;
 Methoxyphenethylamine-CMC;
 Methoxyphenethylamine-HES;
 Arachidonic acid-Hemacell; and
 Arachidonic acid-PEG.
 PE-phosphatidylethanolamine
 PS-phosphatidylserine
 CMC = carboxymethylcellulose (M.W. 25,000)
 HES = hydroxyethyl-starch (M.W. = 40,000)
 PEG = polyethyleneglycol (M.W. 4000)

Cell-impermeable PLA₂ inhibitors of this invention are prepared by linking a carrier, e.g., a polymer, directly or indirectly to a cell-permeable PLA₂ inhibitor according to the general reaction schemes:


- a) inhibitor + spacer \longrightarrow inhibitor-spacer + polymer \longrightarrow inhibitor-spacer-carrier
 b) inhibitor + carrier \longrightarrow inhibitor-carrier

- c) carrier + spacer \longrightarrow carrier-spacer + inhibitor \longrightarrow carrier-spacer-inhibitor
 d) inhibitor + reactive group \longrightarrow inhibitor-reactive group + spacer \longrightarrow inhibitor-reactive group-spacer + carrier \longrightarrow inhibitor-reactive group-spacer-carrier
 e) carrier + reactive group \longrightarrow carrier-reactive group + spacer \longrightarrow carrier-reactive group-spacer + inhibitor \longrightarrow carrier-reactive group-spacer-inhibitor.
 f) inhibitor + reactive group \longrightarrow inhibitor-reactive group;
 carrier + reactive group \longrightarrow carrier-reactive group;
 inhibitor-reactive group + carrier-reactive group + spacer \longrightarrow inhibitor-reactive group-spacer-reactive group-carrier.

With acylated phosphatidylethanolamine (PE) used as precursor for the PLA₂ inhibitor, various lengths of dicarboxylic acids can be used as spacers. These acids can be linked to natural, semi-synthetic or synthetic PE. For example, PE can be linked to aminodextran indirectly as follows:




Ia



IIa


5

10



15

20




25

30

35

40





IIIa

45


Polymers with carboxylic groups, such as polyamino acids, carboxymethyl cellulose or polymers to which fatty acids have been linked, can be linked directly to PE according to the following scheme:




50

55



NBD lyso PS can be linked to an amino polymer according to the following scheme:






Glycerol ether, amine, amide, thioether, ester and thioester derivatives of the general formula:



wherein X is -O-, -S-,

-O-C(=S)-, -O-C(=O)-, -NH-C(=O)-, -NH-; R₁ is alkyl; R₂ is alkyl; and R₃ is alkyl,

-C(=O)-alkyl, phosphoryl-choline, phosphoryl-serine, phosphoryl-ethanolamine, phosphoryl-glycerol, phosphoryl inositol, etc., can be linked to the carrier moiety via the R group according to the following reaction scheme:



Ethylene glycol derivatives of the general formula:



wherein R₁ = alkyl,

10 $\begin{array}{c} \text{O} \\ || \\ \text{C} \end{array}$ -alkyl, phosphoryl-choline, phosphoryl-serine, phosphoryl-ethanolamine, phosphoryl-glycerol, or phosphorylinositol, etc., and X is -O-, -S-,

15 $\begin{array}{c} \text{S} \\ || \\ \text{O} - \text{C} \end{array}$ -, $\begin{array}{c} \text{O} \\ || \\ \text{O} - \text{C} \end{array}$ -, $\begin{array}{c} \text{O} \\ || \\ \text{NH} - \text{C} \end{array}$ -, $\begin{array}{c} \text{O} \\ || \\ \text{NH} - \end{array}$ can be linked to carriers, e.g., by the following reaction scheme, which is an example of the preparation of ethylene glycol monoether phosphatidyl compounds of this invention:

20

25

30

35

40



45

50


55



10




35




50 Derivatives of p-methoxyphenethylamine can be bound to a carrier molecule, e.g., dextramine, according to this invention as follows:

55



I f


10



15

II f


20



25

III f


30



35

IV f

40



V f

45

Uses of PLA₂ Inhibitors of this Invention

50

PLA₂ is present in animal venom, e.g., of snakes and other venomous reptiles, and in the toxin injected by insect stings, e.g., of bees. The hemolysis and neurotoxicity induced by snake bites and insect stings is mediated by the action of this enzyme on cell surface phospholipids. The PLA₂ inhibitors of this invention are therefore useful for the treatment of venom toxicity.

55

The inhibitors of cell surface PLA₂ of this invention can also be used for treatment of diseases associated with excessive endogenous or exogenous PLA₂ activity, e.g., in the form of pharmaceutical compositions adapted for such treatment.

The novel extra-cellular inhibitors of this invention comprise a PLA₂ inhibitor linked directly or via a suitable spacer to a carrier, the latter generally but not necessarily being a high-molecular-weight substance, such as a polymer. The products of the invention act as extra-cellular inhibitors of PLA₂ and thus do not exhibit the cytotoxic effects of PLA₂ inhibitors which penetrate into the cells.

The preferred inhibitors of this invention are phosphalidylethanolamine and phosphatidylserine linked directly to or via a divalent bridging moiety to a carrier.

The novel compositions of the invention are useful for the treatment of a wide variety of PLA₂-associated conditions, as described hereinafter.

The activity of PLA₂ in cell surface membranes has been correlated with cellular secretion in general. Examples of such cell-secreted entities are neurotransmitters, histamine, prostaglandins, leukotrienes, thromboxanes, cholesterol, triglycerides, peroxides, lysosomal enzymes, and secretory products in general. Of special interest are the eicosanoids (prostaglandins, thromboxanes and leukotrienes) which are metabolites of arachidonic acid released from phospholipids by the action of PLA₂. Regulation of PLA₂ in cell surface membranes thus provides a treatment for pathological conditions associated with oversecretion of these substances, such as occurs in allergic response, inflammation, atherosclerosis, thrombosis, myocardial infarct, hypertension, and neurological disorders, among others, and ameliorating the adverse symptoms associated therewith.

The compounds of this invention have been found to possess the valuable pharmacological properties of the PLA₂ inhibitor moiety thereof but lack the side effects thereof associated with cell penetration thereby.

The compounds are particularly useful as anti-inflammatory agents, in the treatment of allergies and oversecretion-related diseases in general. They can be employed in admixture with other drugs indicated for the particular condition being treated.

The compounds of this invention are generally administered to animals, including but not limited to mammals, e.g., livestock, household pets, humans, cattle, cats, dogs, poultry, etc. The pharmacologically active compounds of this invention can be processed in accordance with conventional methods of galenic pharmacy to produce medicinal agents adapted for administration to patients suffering from a PLA₂ associated abnormal condition.

The compounds of this invention can be employed in admixture with conventional excipients, e.g., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral (e.g., oral) or topical application which do not deleteriously react with the active compounds. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohols, gum arabic, vegetable oils, benzyl alcohols, polyethylene glycols, gelatine, carbohydrates such as lactose, amylose or starch, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, hydroxy methylcellulose, polyvinyl pyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds. They can also be combined where desired with other active agents, e.g., vitamins.

For parenteral application, particularly suitable are injectable, sterile solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. Ampoules are convenient unit dosages.

For enteral application, particularly suitable are tablets, dragees, liquids, drops, suppositories, or capsules. A syrup, elixir, or the like can be used when a sweetened vehicle is employed.

Sustained or directed release compositions can be formulated, e.g., liposomes or those wherein the active compound is protected with differentially degradable coatings, e.g., by microencapsulation, multiple coatings, etc. It is also possible to freeze-dry the new compounds and use the lyophilizates obtained, for example, for the preparation of products for injection.

Generally, the compounds of this invention are dispensed in unit dosage form comprising 1 to 1,000 mg per unit dosage in a pharmaceutically acceptable carrier.

The dosage of the compounds according to this invention generally is 0.1 to 100, preferably 1 to 20 mg/kg/day, when administered to patients, e.g., humans, to treat, e.g., subcutaneously to treat arthritis, analogously to the known agent, dexamethasone.

It will be appreciated that the actual preferred amounts of active compound in a specific case will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, and the particular situs and organism being treated. Dosages for a given host can be determined using conventional considerations, e.g., by customary comparison of the differential activities of the subject compounds and of a known agent, e.g., by means of an appropriate, conventional pharmacologi-

cal protocol.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

In the foregoing and in the following examples, all temperatures are set forth uncorrected in degrees Celsius and unless otherwise indicated, all parts and percentages are by weight.

The entire texts of all applications, patents and publications, if any, cited above and below are hereby incorporated by reference.

Example 1: PVA-PE

(a) Preparation of phosphatidylethanolamine (PE) dodecandioic derivative:

300 μ mol of dodecandioic acid were added to 25 μ mol of PE dissolved in 2 ml of dry dichloromethane, followed by 100 mg of dicyclohexyl-carbodiimide (DCC), 0.2 ml of triethylamine and 0.5 ml of absolute methanol. The mixture was incubated for 24 hours at 40°C. Analysis of the products by thin layer chromatography, in chloroform:methanol:acetone:acetic acid:water (3:1:4:0.5), showed two spots sensitive to phosphate spray, identified as PE, near the origin, and Ac-PE at R_f = 0.85. This system was used for purification of Ac-PE on silica gel column.

(b) Binding of PVA to dodecandioyl-PE:

200 mg EDCC (1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide) were added to 10 μ mol of Ac-PS and 50 mg of PVA-hydrazide in 0.5 ml DMSO followed by addition of 200 mg EDCC and stirred for 7 h at 45°C. The PVA conjugate was precipitated and repeatedly washed with ethanol to remove the DMSO and the EDCC. The precipitate was dissolved in 4 ml distilled water, extensively dialyzed against water and lyophilized.

Example 2: Dextranamide-PE

Preparation of dextranamine:

10 g of dextran-40, MW 40,000, were dissolved in 40 ml of water and oxidized with 0.6 g of sodium periodate during 2 h of stirring with a magnetic stirrer. Thereafter, the oxidized dextran was eluted through a column of 100 ml Dowex 1X8-100 (chloride form) with distilled water. The solution of oxidized dextran was mixed with (5 g) diamino hexane for 1 h, then the schiff base formed was reduced by sodium borohydride 50 mmol. After 5 h of stirring, the conjugate was precipitated with ethanol redissolved in distilled H₂O, dialyzed against water and lyophilized.

Preparation of dextranamine dodecanoyl PE:

10 μ mol of acetyl-PE, prepared as in Example 1, were reacted with 50 mg of dextranamine in 100 ml DMSO and 200 mg DCC; the mixture was stirred for 7 h at 45°C and the conjugate was precipitated with EtOH washed with EtOH, then redissolved in distilled H₂O, dialyzed extensively against H₂O, and lyophilized.

Example 3: CMC-Acyl-PE

10 μ mole PE were dissolved in 2 ml DMSO and reacted with 50 μ mol succinic acid in the presence of 500 mg DCC during 2h at 50°C. The carboxyacyl PE formed, was purified on a silica acid column and

reacted with 500 mg diaminoethane in 5 ml of chloroform:methanol:1:1, in the presence of 500 mg DCC. The resulted amino derivative of PE was directly coupled to carboxymethylcellulose (CMC, sigma C-8758), by reacting the acyl-PE with 1 g of CMC in 10 ml H₂O in the presence of 200 mg 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDCC). After an overnight stirring the reaction mixture was dialysed
 5 extensively against H₂O and lyophilized.

Example 4: Dextran-PE

10 Purified carboxy-dextran, prepared by dissolving 10 g of dextran 70 in 40 ml of water and oxidizing and then purifying dextran as described in Example 2, was reacted with 2 g of aminohexanoic acid for 2 hours in H₂O, 100 mole of NaBH₄ were added gradually over a 2-hour period and then left to stir for another 3 h. 2 ml of EtOH were then added and the solution was left to stir another hour. The reaction product then was precipitated from the reaction mixture with EtOH, washed with EtOH, dissolved in H₂O, dialyzed against
 15 H₂O and then lyophilized.

The lyophilized carboxy-dextran was dissolved in DMSO and 50 mg thereof in 2 ml and 10 M of PE were added followed by 100 mg of DCC. After 5 hours incubation at 50 °C, the dextran-PE was precipitated with EtOH, washed with EtOH, dissolved in water, dialyzed and lyophilized.

Example 5: CMC-PE

20 1 gr of carboxymethyl cellulose (low viscosity Sigma C-8758) was dissolved in 50 ml of H₂O. 200 mg of phosphatidyl ethanolamine (PE) was added, followed by 500 mg of 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide. The solution was stirred overnight, dialyzed thoroughly against H₂O and then lyophilized.

Example 6: Poly-D-glutamic acid-PE

30 50 mg poly-D-glutamic acid (MW 50,000-100,000) (Sigma) was reacted with 5 mg of PE in 50 mM phosphate buffer pH 5.2 in the presence of 100 mg of 1-ethyl-3-(3-dimethyl-aminopropyl)carbodiimide. After 5 hours at 45 °C, the solution was dialyzed extensively against distilled water and lyophilized.

Example 7: Polyacrylic acid-PE

35 The reaction was performed as in Example 5, except 50 mg of polyacrylic acid (MW 90,000 or 5,000) were used as the starting polymer.

Example 8: NBD-PS Linked to Dextranamine via Dodecandioic Acid

40 20 mg of lyso PS were reacted with 40 mg of NBDCl in 5 ml of 2:1 MeOH:H₂O. Et₃N was added until pH - 8.5 was reached and the pH was kept at 8.5 therewith with stirring for 5 hours. Then the mixture was evaporated to dryness in a Rotavapour at 20 °C, then applied to a silica gel column, which was eluted with CH₂Cl₂, then successively with CH₂Cl₂:MeOH (9:1, 8:2 and 7:3), then CH₂Cl₂:MeOH:H₂O (75:25:4, 65:35:5 and finally 50:50:8), where the NBD-lyso-PS eluted as an orange band. The solution was evaporated to dryness. The resulted NBD-lyso-PS (10 mg) was reacted with 20 mg of the di-N-hydroxysuccinimide ester of dodecandioic acid in 1 ml DMSO, in the presence of 5 mg dimethylaminopyridine. The resulting orange
 50 compound was purified by preparative TLC on silica gel plates and then reacted with aminodextran, prepared as described in Example 2 in DMSO, by the addition of DCC followed by incubation for 3 hours at 50 °C.

Example 9: Polyethyleneimine-PS

55 50 mg of polyethyleneimine (MW 20,000) were reacted with 10 μM PS in DMSO and 100 mg DCC over a 5-hour period at 45 °C. Thereafter, the solution was dialyzed extensively against H₂O and lyophilized.

Example 10: Hydroxyethylstarch-PE

50 g of hydroxyethylstarch (HES, Sigma H-6382) were dissolved in 0.2 N NaOH 100 ml, and 2 ml of epichlorohydrin were added. The mixture was stirred for 2 hours at 40° and then was added to a solution of 3 g 1,6-diaminohexane in 200 ml of 0.1 N NaOH. The mixture was stirred for 24 hours at 50° and then precipitated with 3 vol of ethanol, redissolved in H₂O, dialyzed against H₂O and then lyophilized. The resulting amino-HES was reacted with acyl-PE as in Example 2.

10 Example 11: Hemacell-linked to 1-dodecanedioate-2-hexadeconyl amidoglyceryl-phosphonylcholine

DL-serine methyl ester (Ig) (1 mol) was added dropwise in 250 ml dry THF to a solution of 0.5 mol LiAlH₄ in 1000 ml of dry THF under N₂ in a 3000 ml round-bottom flask equipped with a magnetic stirrer, a reflux condenser and a dropping funnel. After stirring for 2 h, 100 ml of ethyl acetate were added slowly from the dropping funnel followed by 100 ml methanol, 200 ml H₂O and 200 ml 1N HCl. The THF phase was separated in a separatory funnel and the water phase was extracted 3 times with CH₂Cl₂. The combined THF and CH₂Cl₂ phases were dried on MgSO₄ and evaporated to dryness.

The amino dialcohol (IIg) was acylated by dissolving 0.5 mol of it in 500 ml CH₂Cl₂ in presence of 1.0 mol palmitic acid 2.0 mol Et₃N and 4 mols of dicyclohexylcarbodiimide. After 2 hours reflux the solution was filtered and evaporated to dryness. The amide was purified on silicic acid column eluted with CH₂Cl₂:MOH mixtures.

The dialcohol amide (IIg) formed (0.1 mol) was dissolved in 500 ml of anhydrous benzene containing 0.12 mol Et₃N at 0-5° C. 0.075 mol of 2-chloro-2-oxo-1,3,2-oxaphosphalane were added slowly in 100 ml anhydrous benzene. The mixture was stirred for 12 hours under N₂ at R.T. then filtered and evaporated to dryness.

The phosphate triester thus formed (IVe) was transferred to a pressure bottle, dissolved in 100 ml anhydrous acetonitrile and 0.2 mol of trimethylamine were added. The bottle was sealed and kept at 70° for 24 h, then cooled and the reaction product was collected by filtration.

The thus-produced phosphoryl choline amido alcohol (Vg) 0.05 mol was acylated in 200 ml dry CH₂Cl₂ using 0.1 mol dodecandioic acid and 0.1 mol dimethylaminopyridine and 0.5 mol DCC. The mixture was refluxed for 5 hours then stirred for 2 hours with 100 ml 1 M Na₂CO₃. Then the phases were separated and the water phase was extracted with 4:1 CH₂Cl₂:MeOH. The combined organic phase was washed with H₂O, dried on MgSO₄ and evaporated to dryness.

The resulting oil was purified on a silicic acid column to give Compound VIg, which was reacted with Hemacell by dissolving 10 mmole of VIg in 10 ml H₂O and reacting it with 10 g of Hemacell by addition of 20 g of EDCC for 5h with stirring. The coupled Hemacell derivative dialyzed extensively against H₂O and lyophilized.

40 Example 12: Alginic Acid Derivatives of ethylene glycol phosphorylethanolamine

2.5 g powdered KOH were suspended in 100 ml xylene and 5 mmol dry glycerol were added. The mixture was refluxed in a 250 ml round-bottom flask equipped with a Dean-Stark apparatus, a reflux condenser, a dropping funnel and a magnetic stirrer. After 2 hours of reflux, tetradecyl methane sulfonate (5 mmol) dissolved in 20 ml of xylene was added dropwise. The reflux was continued for 5 hours and then 50 ml of xylene were distilled. The mixture was left to cool, water was added and the reaction product was extracted with CH₂Cl₂ dried on MgSO₄, evaporated to dryness and purified on a silicic acid column.

The ethylene glycol monoether was phosphorylated as in Example II. The phosphate triester formed was opened as in Example 11 using liquid ammonia instead of trimethylamine.

50 The resulting phosphoryl ethanolamine derivative was linked to alginic acid by suspending 1 mmol of the derivatized PE in 100 ml H₂O and reacting it with 5 g. of alginic acid. After 40 min of stirring, 2g. of EDCC were added. The mixture was stirred for 12 h, dialyzed extensively against H₂O and lyophilized.

55 Example 13: Dextran - p-methoxyphenethylamine

0.05 mole of p-methoxyphenethylamine was mixed with 0.05 moles of 6-bromohexanoic acid in 40 ml dioxane. 5 ml of 10% NaOH were added through a dropping funnel over a 15-min. period with thorough

stirring. The mixture was further stirred for 2 hours, then acidified with HCl and then extracted with CH_2Cl_2 . The organic layer was washed with H_2O , dried over MgSO_4 and evaporated to dryness. The product, N-(4-methoxyphenethyl)-aminohexanoic acid (Ih) was crystallized from CH_3CN .

Compound Ih was methylated with formaldehyde by adding it to 30 ml of formalin solution, diluted with 30 ml of methanol. The mixture was stirred for 30 min. then 200 mg of NaBH_4 were added followed by 3 successive additions of 200 mg NaBH_4 every 20 minutes, with continuous stirring. The reaction mixture was stirred overnight and then 50 ml of H_2O and 60 ml of CH_2Cl_2 were added. The organic layer was separated, washed twice with 30 ml of H_2O , dried over MgSO_4 and evaporated to dryness. The resultant product, N-methyl-N-(4-methoxyphenethyl)-aminohexanoic acid (IIh), was esterified with N-hydroxysuccinimide by reacting 0.1 mole of IIh with 0.5 mole DCC and 0.11 moles of N-hydroxysuccinimide in 50 ml of CH_2Cl_2 with stirring for 5 hours at room temperature. The solution was then filtered off and evaporated to dryness. The product, N-hydroxysuccinimide of IIh (IIIh), was crystallized from acetonitrile.

IIIh was conjugated to aminodextran (MW = 4000), prepared as in Example 2, by mixing 20 mg of IIIh, 0.5 g of aminodextran dissolved in 10 ml of H_2O , and 100 mg of ethyldimethylaminopropylcarbodiimide (EDCC). The mixture was stirred overnight and then precipitated, washed with EtOH, redissolved in H_2O and reprecipitated with EtOH. The thus-produced dextran-bound p-methoxyphenethylamine was dissolved in water, dialyzed extensively against H_2O and lyophilized.

Activity Data

The activity of phospholipase- A_2 in the surface membrane of intact cells was determined in various cell types. Among them were endothelial and mast cells, hepatocytes, platelets and oligodendrites. This activity was inhibited with increasing concentration of extracellular inhibitors; (e.g., The activity of PLA_2 in the surface membrane of cultured rat hepatocytes was practically blocked at 100 μM /mg cell protein of bound PS. In parallel, secretion from these cells (e.g., of prostacyclin, thromboxane, histamine, lysosomal enzymes, triglycerides) as well as platelets aggregation was inhibited in correlation with the inhibition of PLA_2 activity.

Hydrolysis of liposomal phospholipids, as well as hemolysis of human red blood cells by snake and bee venom, was inhibited by the PLA_2 inhibitors.

In vivo inhibition of thrombus formation in rabbits was obtained by I.V. administration of PE bound to dextran, CMC or Hemacell, to a final PE concentration in plasma of about 200 μM . The application of the active substance is by way of injection, oral (coated or not), rectal or aerosol. The dosage is generally about 5-50 mg/kg of body weight.

The following examples illustrate the PLA_2 -inhibiting activity of the cell impermeable compounds of this invention. Results obtained with dextranhydrazide-PS (Dex-AC-PS) are given for comparison to demonstrate the efficiency of the compounds of this invention.

Example 14:

Inhibition of Phospholipase A_2 from human PMN leukocytes by cell-impermeable inhibitors (PI) of phospholipase A_2

Method

The test compound is incubated (37°, 60 min., pH 7.0) with crude phospholipase A_2 (extracted from human polymorphonuclear leukocytes), in the presence of 2.5×10^8 autoclaved E.coli containing 5 nmol phospholipids labelled with $[1-^{14}\text{C}]$ -oleate, and 0.5 mmol/l CaCl_2 . The reaction is stopped by extraction with modified Dole reagent. Free oleic acid is separated on a disposable silicic acid (Kieselgel G 100) column and the radioactivity is determined. Percent inhibition is calculated after subtracting a blank (assay mixture without enzyme).

Results**Inhibition of PLA₂ from human PMN**

5	PLI-type	IC ₅₀ (ug/ml)
	a) CMS*	40
10	b) HMS+	57
	c) Dex-Ac-PS*	720
15	*CMS = carboxymethylcellulose-phosphatidylserine	
	+HMS = Hemacell-phosphatidylserine	
	*, Dextran-hydrazide-dodecanedioic-phosphatidylserine	

Example 15**Inhibition of PLA₂-oedema in rats by cell-impermeable inhibitors (PLI) of phospholipase A₂****Method**

By subplantar injection of 200 U PLA₂ (ca. 0.3 mg purified phospholipase A₂ from porcine pancreas, Boehringer Mannheim) in 100 µl physiological saline into the left hind paw of male or female rats (LEW/TIF, ca. 200 g body weight, 5 animals per group) a localized edema was induced and its intensity was determined plethysmometrically 3 and 5 hours later. Test compounds were suspended in 0.75% methylcellulose and administered i.p. (5 ml/kg) 1 hour prior to the PLA₂ injection. The edema-inhibiting effect was expressed as percent inhibition in comparison to vehicle-treated controls.

Results: Inhibition of PLA₂-oedema in rats

	PLI type	Dose (mg/kg)	% Inhibition	
			3 hours	5 hours
5				
10	a) CMS	1 X 20 i.p.	15	29
	b) "	100 i.p.	51	54
15	c) HMS	1 X 20 i.p.	36	62
	d) "	100 i.p.	63	80
20	e) HME*	1 X i.p.	0	0
	f) "	i.p.	27	25
	g) "	i.p.	53	45

25 *HME = Hemacell-phosphatidylethanolamine

30 Example 16

Inhibition of Prostaglandin (PGE₂) and Leukotriens (LTC₄) production in mouse Peritoneal Macrophages by cell-impermeable inhibitors (PLI) of phospholipase A₂ (PLA₂):

35

Method

40 Peritoneal cells from NMRI mice were obtained by lavage with Dulbeccos MEM. The cells were washed and plated at 2×10^6 well in Dulbeccos MEM with FCS in 96-well plates. After 2 hours (or overnight) incubation at 37°C, the adherent macrophages were washed 3 times. The medium was replaced by Dulbeccos MEM with lactalbumin hydrolysate. Test compounds were suspended in water. One hour later, the macrophages were stimulated with 10^{-5} M phorbol-myristate-acetate. Another 2 hours later, PGE₂ and LTC₄ were measured in the supernatants by radioimmunoassay. The results are expressed as IC₅₀ of
 45 PGE₂- and LTC₄-production. (IC₅₀ = PLI concentration which inhibits PGE₂ or LTC₄ production by 50%)

50

55

Results: PLI-IC₅₀ concentration for PGE₂ and LTC₄ production:

	PLI type	IC ₅₀ for	IC ₅₀ for
		PGE ₂ production (µg/ml)	LTC ₄ production (µg/ml)
a)	CMS	100	7
b)	"	100	19
c)	"	> 100	28
d)	HMS	> 100	1
e)	"	> 100	10
f)	Dex-Ac-Ps	300	1000
g)	"	600	400

Example 17

Inhibition of Serotonin secretion and Phospholipase A₂ (PLA₂) activity in Rat Basophilic Leukemia (RBL) cells, by cell-impermeable inhibitor (PLI)

A. Inhibition of Serotonin Secretion with HMS

Inhibition of serotonin secretion by cell-impermeable PLA₂ inhibitor:

Cultured RBL were incubated with tritiated serotonin (10⁷ DPM/ml/10⁶ cell) for 2 hours, then washed. Serotonin secretion was activated in the absence or presence of HMS (1.5 mg/ml) by the addition of the ionophore A23187 (0.3 µM) for the desired time. The radioactive serotonin accumulated in the culture medium during the activation period was determined. In the absence of HMS, approximately 1.3 and 1.9 DPM x 10⁻⁵ was excreted in 15 and 30 minutes, respectively, whereas in the presence of the HMS, only about 0.3 DPM x 10⁵ was secreted in each of these time periods.

B. Inhibition of PLA₂ Activity with HMS

Cultured RBL cells were incubated with the fluorescent substrate of PLA₂ (C₆-NBD-PC) as described by Yedgar et al. (1986) FEBS Letters, 200:165-168. After 1 hour of incubation, the cultures were subjected to lipid extraction and determination of C₆-NBD-PC hydrolysed as previously described.

Results:

HMS	C ₆ -NBD-PC hydrolysed
<u>concentration (mg/ml)</u>	<u>(nmole/mg cell protein-hour)</u>
none (control)	1.42 \pm 0.19
0.5	0.37 \pm 0.04
1.5	0.11 \pm 0.07

Example 18Inhibition of human platelet aggregation and Thromboxane (TXB₂) secretion by cell-impermeable inhibitors (PLI) of phospholipase A₂Method:

Platelet-rich plasma (PRP) was obtained from human donors by centrifugation. Platelet aggregation at 37° C and TXB₂ secretion were induced by the addition of an aggregating agent (inducer) as indicated in the table below. Platelet aggregation was determined by measurement of the suspension optical density in a double beam aggregometer. Results are expressed as % of the control aggregation obtained by the addition of the inducer in the absence of an inhibitor. TXB₂ secretion to the extracellular medium, was determined by radioimmunoassay. When the test PLI was applied, the PRP was incubated with the inhibitor for 10 min at 37° C, prior to induction of aggregation.

**INHIBITION OF PLATELET AGGREGATION AND THROMBOXANE (TXB₂)
SECRETION BY CELL-IMPERMEABLE INHIBITORS (PLI) OF PHOSPHOLIPASE
-A₂ (PLA₂)**

PLI type	Aggregating agent	PLI conc. (mg/ml)	Aggregation (% of control)	TXB ₂ secreted (ng/ml)
HME [†]	ADP (2 µg/ml)	none	100	82±4
"	"	0.70	6	14±7
"	Collagen (40 µg/ml)	none	100	42±6
"	"	0.85	13	17±7
"	Epinephrine (0.8 µg/ml)	none	100	
"	"	0.75	9	
ALS ^{**}	Epinephrine (0.8 µg/ml)	none	100	
"	"	1.5	5	
CME ⁺	Epinephrine (0.8 µg/ml)	none	100	
"	"	1.5	0	
CMS ⁺⁺	Epinephrine (0.8 µg/ml)	none	100	
"	"	1.5	0	
ALZ [*]	Epinephrine (0.8 µg/ml)	none	100	
"	"	0.7	9	
HMS ^{††}	Epinephrine (0.8 µg/ml)	none	100	
"	"	1.2	2	
Dex-A ₂ -PS	ADP [*] (2 µg/ml)	3.0	14	
"	Epinephrine (0.8 µg/ml)	6.5	19	

^{*}, alginic acid - phosphatidylethanolamine

^{**}, alginic acid - phosphatidylserine

⁺, carboxymethylcellulose - phosphatidylethanolamine

⁺⁺, carboxymethylcellulose - phosphatidylserine

⁼, Hemacell - phosphatidylethanolamine

⁼⁼, Dextran-hydrazide-dodecanedioyl-phosphatidylserine

Example 19 - Inhibition of prostacyclin production in endothelial cell by cell-impermeable PLA₂ inhibitors (CMS and HME)

Method:

Cultured adrenal capillary endothelial cells were stimulated to produce prostacyclin (6-keto-PGF_{1α}) by bradykinin in the absence and presence of CMS or HME. The prostacyclin, secreted to the culture medium was determined by radioimmunoassay.

EFFECT OF CELL-IMPERMEABLE PLA₂ INHIBITORS ON PROSTACYCLIN PRODUCTION BY ENDOTHELIAL CELLS

Treatment	6-keto-PGF _{1α} PROD. (PG/10 ⁻⁶ Cells)	% Inhibition
Bradykinin (1 μM)	201.9 ± 21.6	
CMS (1 mg/ml)		
+ Bradykinin (1 μM)	78.7 ± 6.8	61
HME (1mg/ml)		
+ Bradykinin (1 μM)	36.4 ± 7.2	82

Example 20 - Antiinflammatory activity of cell-impermeable PLA₂ inhibitors

Two compounds (CMS and MHS) were tested in the adjuvant arthritis model of inflammation. Paw swelling is induced by an injection of Freund's adjuvant into the right hind paw. Systemic inflammation spreads to the contralateral paw within 2 weeks. The compounds were administered subcutaneously from day 3 to day 14 at a dose of 70 mg/kg. By day 14, paw diameter (measured with calipers) was significantly decreased in arthritic animals dosed in HMS and CMS. The noninjected and injected paws of arthritic rats treated with HMS exhibited a decrease in paw swelling of 43% (P ≤ 0.01) and 19% (P ≤ 0.01), respectively. CMS decreased noninjected paw swelling 51% (P ≤ 0.005) and injected paw swelling 30% (P ≤ 0.001). There were no gross signs of toxicity over the two week dosing period. No drug treated group lost weight compared to the arthritic controls.

The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

Claims

1. A compound capable of inhibiting the enzyme phospholipase A₂ (PLA₂) comprising a cell-permeable PLA₂-inhibitor moiety covalently bonded directly or indirectly to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, with the proviso that phosphatidylserine is not bonded indirectly via a divalent didodecanedioyl group to dextran-hydrazide.

2. A compound as claimed in claim 1 wherein the carrier moiety is other than dextran-hydrazide.

3. A compound as claimed in claim 1 or 2 wherein the PLA₂-inhibitor moiety is phosphatidyl-ethanolamine or phosphatidyl-serine.
4. A compound as claimed in any of claims 1 to 3, wherein the PLA₂-inhibitor moiety is an acylated phosphatidyl-ethanolamine or acylated phosphatidyl-serine.
- 5 5. A compound as claimed in any of claims 1 to 4, wherein the carrier moiety is a polymer.
6. A compound as claimed in any of claims 1 to 5, wherein the carrier moiety is a degraded gelatin polypeptide cross-linked via urea bridges.
7. A compound as claimed in any of claims 1 to 6, wherein the carrier moiety is carboxymethylcellulose.
- 10 8. A compound as claimed in any of claims 1 to 7, wherein the carrier moiety is alginic acid, hydroxyethyl starch polyethylene glycol or dextran.
9. A compound as claimed in any of claims 1 to 8, wherein the PLA₂-inhibitor moiety is separated from the carrier moiety by a divalent bridging moiety.
10. A compound capable of inhibiting the enzyme phospholipase A₂ (PLA₂) comprising a cell-permeable PLA₂-inhibitor moiety covalently bonded directly or indirectly to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, with the proviso that phosphatidylserine is not bonded indirectly via a divalent didodecanedioyl group to dextran hydrazide; for use in medicine.
- 15 11. The use of a compound capable of inhibiting the enzyme phospholipase A₂ (PLA₂) comprising a cell-permeable PLA₂-inhibitor moiety covalently bonded directly or indirectly to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, in the preparation of a PLA₂ inhibiting agent.
12. A pharmaceutical composition comprising a compound capable of inhibiting the enzyme phospholipase A₂ (PLA₂) comprising a cell-permeable PLA₂-inhibitor moiety covalently bonded directly or indirectly to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, with the proviso that phosphatidylserine is not bonded indirectly via a divalent didodecanedioyl group to dextran-hydrazide.

30 Claims for the following Contracting States: ES, GR

1. A process for the preparation of a compound capable of inhibiting the enzyme phospholipase A₂ - (PLA₂), the process comprising covalently bonding directly or indirectly a cell-permeable PLA₂-inhibitor moiety to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, with the proviso that phosphatidylserine is not bonded indirectly via divalent didodecanedioyl to dextran-hydrazide.
2. A process as claimed in claim 1, wherein the carrier moiety is other than dextran-hydrazide.
3. A process as claimed in claim 1 or 2, wherein the PLA₂-inhibitor moiety is phosphatidyl-ethanolamine or phosphatidyl-serine.
- 40 4. A process as claimed in any of claims 1 to 3, wherein the PLA₂-inhibitor moiety is an acylated phosphatidyl-ethanolamine or acylated phosphatidyl-serine.
5. A process as claimed in any of claims 1 to 4, wherein the carrier moiety is a polymer.
6. A process as claimed in any of claims 1 to 5, wherein the carrier moiety is a degraded gelatin polypeptide cross-linked via urea bridges.
- 45 7. A process as claimed in any of claims 1 to 6, wherein the carrier moiety is carboxymethylcellulose.
8. A process as claimed in any of claims 1 to 7, wherein the carrier moiety is alginic acid, hydroxyethyl starch polyethylene glycol or dextran.
9. A process as claimed in any of claims 1 to 8, wherein the PLA₂-inhibitor moiety is separated from the carrier moiety by a divalent bridging moiety.
- 50 10. A process for the preparation of a pharmaceutical composition, the process comprising admixing a compound capable of inhibiting the enzyme phospholipase A₂ (PLA₂) comprising a cell-permeable PLA₂-inhibiting moiety covalently bonded directly or indirectly to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, with the proviso that phosphatidylserine is not bonded indirectly via a divalent didodecanedioyl group to dextran-hydrazide and a pharmaceutically acceptable carrier.
- 55

11. The use of a compound capable of inhibiting the enzyme phospholipase A₂ (PLA₂) comprising a cell-permeable PLA₂-inhibitor moiety covalently bonded directly or indirectly to a physiologically acceptable carrier moiety capable of inhibiting cell internalisation of the cell-permeable PLA₂-inhibitor moiety, in the preparation of a PLA₂ inhibiting agent.

5

10

15

20

25

30

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 88 30 9996

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
D,A	FEBS LETTERS, vol. 200, no. 1, May 1986, pages 165-168, Federation of European Biochemical Societies; S. YEDGAR et al.: "Synthesis of cell-impermeable inhibitor of phospholipase A2" * Page 166, paragraph 2.1.3; page 166, "results and discussions" *	1,3-4,6	A 61 K 47/00
Y	IDEM ---	2,5	
Y	JOURNAL OF MEDICINAL CHEMISTRY, vol. 28, no. 1, 1985, pages 137-140, American Chemical Society, US; E. HURWITZ et al.: "The covalent linking of two nucleotide analogues to antibodies" * Page 138, scheme I *	2,5	
D,X	BIOCHEMISTRY, vol. 21, no. 6, 1982, pages 1254-1260, American Chemical Society, US; T.W. MARTIN et al.: "Rat mast cell phospholipase A2: activity toward exogenous phosphatidylserine and inhibition by N-(7-nitro-2,1,3-benzoxadiazol-4-yl)phosphatidylserine" * Whole article *	1-3	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 09-02-1989	Examiner BERTE M.J.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	

EPO FORM 1503 03.82 (P0401)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.